

Study of the coincidences between the gravitational wave detectors EXPLORER and NAUTILUS in 2001

P Astone¹, D Babusci², M Bassan³, P Bonifazi⁴, P Carelli⁵, G Cavallari⁶,
E Coccia³, C Cosmelli⁷, S D'Antonio³, V Fafone², G Federici⁷, S Frasca⁷,
G Giordano², A Marini², Y Minenkov³, I Modena³, G Modestino²,
A Moleti³, G V Pallottino⁷, G Pizzella⁸, L Quintieri², A Rocchi³,
F Ronga², R Terenzi⁴, G Torrioli⁹ and M Visco⁴

¹ Istituto Nazionale di Fisica Nucleare INFN, Rome, Italy

² Istituto Nazionale di Fisica Nucleare INFN, Frascati, Italy

³ University of Rome 'Tor Vergata' and INFN, Rome, Italy

⁴ IFSI-CNR and INFN, Rome, Italy

⁵ University of L'Aquila and INFN, Rome, Italy

⁶ CERN, Geneva, Switzerland

⁷ University of Rome 'La Sapienza' and INFN, Rome, Italy

⁸ University of Rome 'Tor Vergata' and INFN, Frascati, Italy

⁹ IESS-CNR, Rome, Italy

Received 23 July 2002, in final form 24 September 2002

Published 16 October 2002

Online at stacks.iop.org/CQG/19/5449

Abstract

We report the result from a search for bursts of gravitational waves using data collected by the cryogenic resonant detectors EXPLORER and NAUTILUS during 2001 for a total measuring time of 90 days. With these data we repeated the coincidence search performed on the 1998 data (which showed a small coincidence excess) applying data analysis algorithms based on known physical characteristics of the detectors. With the 2001 data, a new interesting coincidence excess is found when the detectors are favourably oriented with respect to the galactic disc.

PACS numbers: 0480, 0430

1. Introduction

Cryogenic gravitational wave (GW) antennas entered into a long-term data taking operation in 1990 (EXPLORER [1]), 1991 (ALLEGRO [2]), 1993 (NIOBE [3]), 1994 (NAUTILUS [4]) and in 1997 (AURIGA [5]), with gradual performance improvements over the years.

Analysis of the data taken in coincidence among all cryogenic resonant detectors in the operation during 1997 and 1998 was performed [6]. No coincidence excess was found above background using the event lists produced under the protocol of the International

Table 1. Main characteristics of the two detectors in 2001. The axes of the two detectors are aligned to within a few degrees of one another, the chance of coincidence detection being thus maximized. The pulse sensitivity for both detectors is of the order of $h \sim 4 \times 10^{-19}$ for 1 ms bursts.

Detector	Latitude	Longitude	Azimuth	Mass (kg)	Frequencies (Hz)	Temperature (K)	Bandwidth (Hz)
EXPLORER	46.45 N	6.20 E	39° E	2270	904.7, 921.3	2.6	~9
NAUTILUS	41.82 N	12.67 E	44° E	2270	906.97, 922.46	1.5	~0.4

Gravitational Event Collaboration (IGEC) among the groups ALLEGRO, AURIGA, EXPLORER/NAUTILUS and NIOBE.

Later [7], a coincidence search between the data of EXPLORER and NAUTILUS was carried out by introducing analysis considerations based on physical characteristics of the detectors: the event energy and the directionality. The result was a small coincidence excess when the detectors were favourably oriented with respect to the galactic centre.

Here we extend our analysis to new data obtained in 2001, when both EXPLORER and NAUTILUS were operating at their best sensitivity, using the same procedures applied to the previous analysis [7]. As in [7], we will sometimes use the word *probability*, although we are well aware that its significance might be jeopardized by any possible data selection. With this *proviso* we will use probability estimations in comparing different experimental conditions.

2. Experimental data

The resonant mass GW detectors NAUTILUS, operating at the INFN Frascati Laboratory, and EXPLORER, operating at CERN, both consist of an aluminium bar with a mass of 2270 kg cooled to very low temperatures. A resonant transducer converts the mechanical oscillations into an electrical signal and is followed by a dc-SQUID electronic amplifier. The bar and the resonant transducer form a coupled oscillator system with two resonant modes.

With respect to 1998, the following changes were made in the set-up of the detectors: the NAUTILUS detector operated at a thermodynamic temperature of 1.5 K instead of 0.14 K, and EXPLORER was equipped with a new transducer providing a larger bandwidth and consequently enhanced sensitivity. The characteristics of the two detectors are given in the table 1.

The data, sampled at intervals of 12.8 ms for NAUTILUS and 6.4 ms for EXPLORER, are filtered with an adaptive filter matched to delta-like signals for the detection of short bursts [8]. This search for bursts is suitable for any transient GW that shows a nearly flat Fourier spectrum at the two resonant frequencies of each detector. The metric perturbation h can either be a millisecond pulse, a signal made by a few millisecond cycles or a signal sweeping in frequency through the detector resonances. This search is, therefore, sensitive to different kinds of GW sources, such as a stellar gravitational collapse, the last stable orbits of an inspiraling neutron star or black-hole binary, its merging and its final ringdown.

Let $x(t)$ be the filtered output of the detector. This quantity is normalized, using the detector calibration, such that its square gives the energy innovation of the oscillation for each sample, expressed in kelvin units.

For well-behaved noise due only to the thermal motion of the oscillators and to the electronic noise of the amplifier, the distribution of $x(t)$ is normal with zero mean. Its variance




Figure 1. Differential probability that the event has the signal-to-noise ratio shown on the abscissa when the signal has $R_s = 20$ (near the threshold $R_t = 19.5$) and $R_s = 30$.

(average value of the square of $x(t)$) is called *effective temperature* and is indicated by T_{eff} . The distribution of $x(t)$ is

$$f(x) = \frac{1}{\sqrt{2\pi T_{\text{eff}}}} e^{-\frac{x^2}{2T_{\text{eff}}}}. \quad (1)$$

In order to extract *events* to be analysed from the filtered data sequence, we set a threshold in terms of a critical ratio defined by

$$\text{CR} = \frac{|x| - \overline{|x|}}{\sigma(|x|)} \quad (2)$$

where $\sigma(|x|)$ is the standard deviation of $|x|$ and $\overline{|x|}$ the moving average, computed over the preceding 10 min.

The threshold is set at $\text{CR} = 6$ in order to obtain, in the presence of thermal and electronic noise alone, a reasonable number of events per day (see [7]). This threshold corresponds to energy $E_t = 19.5 T_{\text{eff}}$. When $|x|$ goes above the threshold, its time behaviour is considered until it falls back below the threshold for longer than 3 s. The maximum amplitude and its occurrence time define the *event*.

The searched *events* are those due to a combination of a GW signal of energy E_s and the noise. The theoretical probability of detecting a signal with a given signal-to-noise ratio $R_s = \frac{E_s}{T_{\text{eff}}}$ in the presence of a well-behaved Gaussian noise is [9]

$$\text{probability}(R_s) = \int_{R_t}^{\infty} \frac{1}{\sqrt{2\pi R_e}} e^{-\frac{(R_s+R_e)}{2}} \cosh(\sqrt{R_e \cdot R_s}) dR_e, \quad (3)$$

where R_e is the signal-to-noise ratio for the event and $R_t = \frac{E_t}{T_{\text{eff}}} = 19.5$ for the EXPLORER and NAUTILUS detectors.

The behaviour of the integrand is shown in figure 1. This figure shows the spread of the event energy due to noise for a given R_s of the signal. It shows that signals with $R_s = 20$ (near the threshold) have a probability of not being detected of about 50%, and signals with $R_s = 30$, rather larger than the threshold, still have a probability of not being detected of nearly 15%. The distinction between the two concepts, *signal* and *event*, is essential for our analysis, as discussed in [7].




Figure 2. The distributions of the hourly averages of T_{eff} in kelvin units for EXPLORER and NAUTILUS. We accept only the time periods with hourly averages $\overline{T_{\text{eff}}} \leq 10$ mK.

Computation of the GW amplitude h from the energy signal E_s requires a model for the signal shape. A conventionally chosen shape is a short pulse lasting a time of τ_g , resulting (for optimal orientation, see later) in the relationship

$$h = \frac{1}{4Lf^2} \frac{1}{\tau_g} \sqrt{\frac{kE_s}{M}}, \quad (4)$$

where f is the resonance frequency, L and M are the length and the mass of the bar, respectively, and τ_g is conventionally assumed to be equal to 1 ms (for instance, for $E_s = 1$ mK we have $h = 2.5 \times 10^{-19}$).

3. Data selection

All events which are in coincidence within a time window of ± 5 s with signals observed by a seismometer are eliminated. This criterion cuts out about 8% of the events.

It is observed that the experimental data are affected by non-Gaussian noise which, in some cases, cannot be observed with any other auxiliary detector. Thus, a strategy is needed to select periods during which the detectors operate in a satisfactory way, as discussed in the following paragraphs.

To this end we consider the quantity T_{eff} , which we use in two ways. The first is to compute T_{eff} by averaging x^2 over 1 h of continuous measurements ($\overline{T_{\text{eff}}}$), and the second to consider the T_{eff} averaged during the 10 min preceding each event. For the hourly averages we show the distribution in figure 2. On observing this figure we decided to consider for the coincidence search only the time periods with hourly averages of less than 10 mK.




Figure 3. The distributions of the 10 min averages of T_{eff} , before each event, in kelvin units for the EXPLORER and NAUTILUS events (with hourly $\overline{T_{\text{eff}}} \leq 10$ mK). We accept only the events with the 10 min average $T_{\text{eff}} < 7$ mK. Note that the EXPLORER events are more numerous than the NAUTILUS events because of the different bandwidth.

The distributions for the T_{eff} averaged over the 10 min preceding each event are shown in figure 3 for EXPLORER and NAUTILUS. It should be noted that the number of events is larger for EXPLORER than for NAUTILUS. This depends on the bandwidth Δf which is larger for EXPLORER (see table 1).

On observing these distributions we decided, as a conservative *a priori* data selection, to make a cut and accept only the events for which the corresponding T_{eff} was below 7 mK (and there was no seismometer veto). This meant regarding the bump at 10 mK for NAUTILUS as due to extra noise. We recall that in the previous search [7] with the noisier 1998 data, the cuts on T_{eff} were made at between 25 and 100 mK.

From our previous experience we have found that the detectors operate in a more stationary way when the noise temperature remains low for longer periods of time because this indicates a smaller contribution of extra noise. To make a quantitative check on this we classified the data stretches in various categories according to the length of the continuous periods having hourly averages $\overline{T_{\text{eff}}} \leq 10$ mK, obtaining the figures shown in table 2. From this table we clearly see that the longer the time period of continuous operation with low noise, the smaller the number of events associated with a noise $T_{\text{eff}} \geq 7$ mK.

Finally, in figure 4, we show the distribution of the event energies selected according to $\overline{T_{\text{eff}}} \leq 10$ mK and $T_{\text{eff}} \leq 7$ mK, for each event, belonging to periods with duration ≥ 1 h. We note that, in spite of our selection criteria, we still have several events with large energy, which indicates the presence of extra noise, in addition to the thermal and electronic ones. The only way to eliminate this noise is by means of the coincidence technique.




Figure 4. The distributions of the EXPLORER and NAUTILUS event energies in kelvin units.

Table 2. In the first column we indicate the minimum time length of continuous operation and in the second column, the total common time of measurement. The last columns indicate the number of events, only those during the common time of operation, with hourly $\overline{T_{\text{eff}}} \leq 10$ mK and the percentage of the events having the 10 min average $T_{\text{eff}} > 7$ mK.

Time length	Hours	EXPLORER		NAUTILUS	
		Events	%	Events	%
≥ 1 h	2156	54 762	5.9	11 252	37
≥ 3 h	2082	52 683	5.0	10 887	34
≥ 6 h	1927	50 344	4.1	9 939	31
≥ 12 h	1490	40 105	3.2	7 268	27

4. Searching for coincidences

It is important to establish the time window for the coincidence search. Using simulated signals and real noise, we characterized [10] the dispersion of the time of the event around the time when the signal is applied.

The standard deviation of the time dispersion for a given detector is

$$\sigma_d = \text{const} \frac{1}{\Delta f} \frac{1}{\sqrt{R_e}} \quad (5)$$

for delta signals, where Δf is the detector bandwidth and the $\text{const} = 0.28$ is determined, for the EXPLORER and NAUTILUS detectors, by means of simulation [11]. For a coincidence analysis with two detectors we have

$$\sigma_w = \sqrt{\sigma_{\text{expl}}^2 + \sigma_{\text{naut}}^2} \quad (6)$$

We decided to take as coincidence window $w = \pm 3\sigma_w$, which is the most recent choice of the IGEC collaboration. Since each event has its own σ_w the value of w will be different for each coincidence. Note that the value of w is almost entirely due to the NAUTILUS detector, since EXPLORER has a much larger bandwidth; it turns out to be of the order of $|w| \sim 0.5$ s, about one half of that used in previous searches for coincidences. With the use of $3\sigma_w$ we also take into account the uncertainty of the detector bandwidth and of the simulation procedure.

Analysis in a coincidence search consists of comparing the detected number of coincidences at zero time delay ($\pm w$) with the background, that is, with coincidences occurring by chance. In order to measure the background due to the accidental coincidences, using a procedure adopted since the beginning of the gravitational wave experiments [12], one shifts the time of occurrence of the events of one of the two detectors a number of times. We shifted this 100 times in steps of $\Delta t = 2$ s (uncorrelated data) from -100 s to $+100$ s. For each time shift we get a number of (shifted) coincidences. If the time shift is zero, we get the number n_c of *observed* coincidences. The accidental background is calculated from the average number of the n_{shift} shifted coincidences obtained from the 100 time shifts,

$$\bar{n} = \frac{\sum_{j=1}^{100} n_{\text{shift}}(j)}{100}. \quad (7)$$

This experimental procedure for evaluation of the background has the benefit of handling the problems arising when the distribution of the events is not stationary (see [13]), although this is not the case with the present 2001 data.

5. Energy filter

It is clear that if a coincidence between the two detectors is due to the arrival of a GW burst, we expect the energies of the two coincident events to be correlated, and we can disregard all coincidences whose corresponding event energies are very different, according to the considerations illustrated in figure 1. Thus, we can apply an energy filter with the aim of reducing the background.

The procedure for application of such an energy filter was set in our previous search for coincidences [7]. We considered signals of various energies E_s . For each coincidence found we calculated the R_s for each of the above signal energy E_s using the known values of the (local) T_{eff} of the two events. We then verified whether the two R_e , for the two events of that coincidence, fell within the interval $R_s \pm \Delta R_s$, such that the two limits $R_s - \Delta R_s$ and $R_s + \Delta R_s$ delimitate (see figure 1) an area of 68% (about one standard deviation for well-behaved noise) for a given value of E_s ; that is, we verified the compatibility of the two events. We followed the same procedure for the shifted coincidences in order to estimate the background after application of the energy filter. In this way we reduced, for the 1998 data, the average number of accidental coincidences from $\bar{n} = 223$ to $\bar{n} = 51$.

This procedure is useful, in particular, if the two detectors have different sensitivity, as in the case of the 1998 data, and, consequently, the event thresholds and the event energies are also different. In the case of two detectors with comparable sensitivity, however, one could also consider comparing directly the energies of the coincident events. For the 2001 data, although in this year the two detectors had comparable sensitivities (see figure 2), we decided not to change the procedure used for the 1998 data. We considered GW signals of energy E_s , in a range covering the energies of our events, i.e. E_s from 5 mK to 1 K in steps of 5 mK, and accepted the coincidence (at zero delay or at a shifted time) if the two events fell within the above interval $R_s \pm \Delta R_s$.

6. Sidereal time distribution

In our previous search for coincidences [7], we took into consideration the non-isotropic response of the detector to a GW burst. We had reasoned that, since extragalactic GW signals could not be detected with the present detectors, possible sources should be located in our galaxy, or in the local group. If any of these sources exist, we should expect a more favourable condition of detection when the detectors are oriented with their axes perpendicular to the direction of the potential source, since the bar cross-section is proportional to $\sin^4(\theta)$, where θ is the angle between the detector axis and the direction of the line joining it with the source. We did find a small coincidence excess when angle θ with respect to the galactic centre was larger than a certain value (see figure 3 of [7]). The inconvenience of this method is that, due to the poor statistics, the result has to be presented in an integral type graph, which makes it difficult to appreciate the real statistical significance of the data. Furthermore, hypotheses must be made on the location of the GW source.

In the present search for coincidences, we extend the previous analysis as follows. We still make use of the directional property of the antenna cross-section. As the Earth rotates around its axis, during the day the detector happens to be variably oriented with respect to a given source at an unknown location. Thus, we expect the signal to be modulated during the day; more precisely the modulation is expected to have a period of one sidereal day (with one or more maxima) (see [14, 15]), since the GW sources, if any, are certainly located far outside our solar system.

The principal, key analysis is carried out with the events in the time periods of at least 12 h of continuous data taking (see table 2) to which the energy filter is applied.

Twenty-four categories of events are considered, one per sidereal hour, the sidereal time referred to as position and orientation halfway between EXPLORER and NAUTILUS (this determines the zero local sidereal time which is not essential for the following considerations). Each category includes coincidences totally independent of those in the other categories. For each category in figure 5, we report the number n_c of observed coincidences, the average number \bar{n} of accidental coincidences obtained by using the time shifting procedure and, given \bar{n} , the probability p that a number $\geq n_c$ of coincidences could have occurred by chance. For comparison we also show a histogram produced with the same procedure using solar hours.

One notes a coincidence excess from sidereal hour 3 to sidereal hour 5, which appears to have some statistical significance, as the two largest excesses occur in two neighbouring hours (the events in each hour are totally independent of those in a different hour). We have $n_c = 7$ coincidences in this 2-h interval and $\bar{n} = 1.7$. In contrast, no significant coincidence excess appears at any solar hour.

The accidental coincidences always have a Poissonian distribution. To check this, we have considered for all the above events the accidental coincidences obtained with 10 000 trials, by time shifting from $-10\,000$ s to $+10\,000$ s in steps of 2 s. The distribution of the number of accidental coincidences is shown in figure 6. The agreement between experimental and expected distributions is excellent.

We repeat the analysis for the events belonging to the larger set of continuous data taking lasting 1 h or more (first line of table 2). We obtain the result shown in figure 7. We note that in the sidereal 2 h interval the number of coincident events has increased from seven to eight and the background has become $\bar{n} = 2.6$.

We now proceed to perform a test, to check if this result is compatible with simultaneous physical excitation of the two detectors of possible non-terrestrial origin. We compare the energies of the coincident events: if the events in EXPLORER and NAUTILUS are due to the same cause, we expect their energies to be correlated. This test must be done without




Figure 5. Result with events in the long time periods (≥ 12 h) of continuous operation. The upper graph on the left shows the number of coincidences n_c indicated with the asterisks and the average number \bar{n} of accidentals versus the sidereal hour. The lower graph on the left shows the Poisson probability of obtaining a number of coincidences greater than or equal to n_c . The two graphs on the right show the result using the solar time in hours. We remark that the data points refer to independent sets of events.




Figure 6. The distribution of the number of accidental coincidences obtained with 10 000 trials, with average number $\bar{n} = 25.32$. The asterisks indicate the experimental distribution, and the continuous line indicates the expected Poissonian distribution.

applying the energy filter. Using the events in the time periods with duration ≥ 12 h and without applying the energy filter, we still get seven coincidences. If we consider the time periods with duration ≥ 1 h, we get eight coincidences. The event energies are very strongly correlated, as shown in figure 8. We also studied the energy correlation of the events of the accidental coincidences and found no correlation.

We also performed a coincidence data analysis when no energy filter was applied at all, obviously expecting a larger accidental background. The result is shown in figure 9. We find




Figure 7. Result with events in time periods ≥ 1 h of continuous operation, as in figure 5.




Figure 8. Correlation between the event energies of NAUTILUS with those of EXPLORER for the eight coincidences that occurred in the sidereal hour interval 3 to 5, in time periods ≥ 1 h. The correlation coefficient is 0.96. No energy filter was applied.

that the coincidence excess in the time interval 3 to 5 sidereal hours still shows up, although less clearly, as expected.

A different way of presenting these data is shown in figure 10. This figure shows that at sidereal hours outside the interval 3 to 5 h, the event energies are not correlated. Also it shows the particular behaviour in the 3 to 5 h interval, when the energies of all coincident events are correlated.

The eight events in the 3 to 5 h period are listed in table 3. We have verified that, using the cosmic ray detector of NAUTILUS, these events are not due to cosmic ray showers.

7. Comparison with the 1998 data

The analysis presented here differs slightly (e.g., for the use of the sidereal time) from that applied previously [7]. We therefore present also the 1998 data in terms of the sidereal




Figure 9. Same as figure 5, with no application of the energy filter. The two graphs on the left refer to events in time periods with duration ≥ 12 h. The two graphs on the right refer to events in time periods with duration ≥ 1 h.




Figure 10. Ratios of the energies (NAUTILUS/EXPLORER) for the events in coincidence belonging to periods with duration ≥ 1 h (no application of the energy filter) versus the sidereal hour.

time. We must consider that the 1998 data are noisier than the 2001 data. In particular, the EXPLORER data have a noise T_{eff} ten times larger than that of the 2001 data, whilst the NAUTILUS noise was of the same order.

For the coincidence search, we change the window from $w = \pm 1$ s (the IGEC choice at that time) used in paper [7] to the present $w = \pm 3\sigma_w$ used here. The result is given in figure 11. An effect similar to the one found for the 2001 data is noticed, although weaker than that obtained with the less noisy 2001 data.




Figure 11. The 1998 data with energy filter. The upper graph on the left shows the number of coincidences n_c indicated by asterisks and the average number \bar{n} of accidentals versus the sidereal hour. The lower graph on the left shows the Poisson probability of obtaining a number of coincidences greater than or equal to n_c . The two graphs on the right show the result using the solar time in hours.

Table 3. List of the coincident events at sidereal hours between 3 and 5. No energy filter has been applied. δt is the time difference between the two coincident events and E is the event energy.

day	hour	min	sec	δt (s)	EXPLORER (mK)		NAUTILUS (mK)		Sidereal hour
					E	T_{eff}	E	T_{eff}	
112	13	59	33.26	0.01	94	3.9	128	2.9	4.6
130	11	39	28.07	-0.08	179	3.6	195	5.9	3.4
133	12	35	45.40	-0.12	194	7.0	226	5.7	4.5
166	10	48	6.04	0.39	73	3.2	64	3.0	4.9
198	7	46	40.32	0.43	102	2.9	133	3.6	4.0
278	2	12	29.65	0.37	63	2.9	57	2.6	3.7
296	0	29	40.59	-0.12	96	2.6	130	5.6	3.1
296	1	24	10.46	0.00	87	2.8	93	4.1	4.1

8. Robustness of the statistical analysis

In any data analysis care must be taken to avoid choosing procedures which favour particular results. Thus, we considered the possibility that our result be biased, although involuntarily, by any such choices. In the present analysis, all choices were made *a priori* and already published in the scientific literature; in particular, the IGEC choice for the coincidence window of $\pm 3\sigma_w$, the energy filter (see [1, 7]) and the threshold for definition of an event (see [6]).

Nevertheless, we tested whether our present choices were indeed, by accident, most apt to produce the coincidence excess. This proved not to be the case.

To find the *most favourable* parameters, that is the threshold and the energy filter parameter, we considered only the coincidences in the sidereal hour interval 3 to 5 and minimized the probability of a coincidence excess by chance. We found that the most favourable threshold for the definition of an event is at $R_t = 20.5$, instead of 19.5. The most favourable parameter for the energy filter is 50% instead of 68%.




Figure 12. The 2001 data with *a posteriori* choices (see text). The two graphs on the left (periods ≥ 1 h) are as in figure 7 and those on the right (periods ≥ 12 h) are as in figure 5. Again we remark that the data points in the figure at different sidereal hours are independent of each other.

We thought it interesting to report the result for these most favourable choices, $R_t = 20.5$ for the threshold and 50% for the energy filter. The result is shown in figure 12. We note an indication that the coincidence excess might extend to the sidereal hour interval 3 to 6, including two more coincidences in the period 5 to 6 sidereal hours (see also figure 10). We want to remark that this hour was not included in the optimization process.

As far as the coincidence window is concerned, we found (*a posteriori*) that the best choice for having a coincidence excess in the 3 to 5 sidereal hour interval is $\pm 3.5\sigma_w$, and any coincidence window from $\pm 2.5\sigma_w$ to $\pm 4\sigma_w$ gives comparable results.

9. Discussion and conclusions

The IGEC search for coincidences [6] was performed without applying the event energy algorithms based on the event amplitude and on the directional properties of the detectors; it gave a no coincidence excess. With new data taken in the year 2001 and with improved sensitivity, we repeated the coincidence search with the detectors EXPLORER and NAUTILUS (no other detector was in operation during 2001), applying data analysis algorithms based on known physical characteristics of the detectors, namely, the energy of the events and directionality of the detectors. We obtained a coincidence excess at sidereal hours between 3 and 5.

At a given sidereal time the intersection of the celestial sphere with the plane perpendicular to the detector axis is a circle. We show in figure 13 two of these circles (i.e., at 1 and 13 sidereal hours) in the right ascension–declination plane. In figure 13, the line which indicates the point sources perpendicular to the detector axis (we call this the *line of maximum sensitivity*) moves to the right with sidereal time. This line intersects the location of the galactic centre twice a day (at 4.3 and at 13.6 sidereal hours). Only once per day (at 4.3 sidereal hour) this line overlaps with the entire galactic disc. The overlapping takes place because of the particular orientation of the detectors on the Earth's surface (see table 1), and would not occur for a different azimuth angle of detector orientation.




Figure 13. The position of the galactic disc and the loci of point sources perpendicular to the detector axis at 1 and 13 sidereal hours on the right ascension–declination plane. At sidereal hour 4.3 the locus tends to coincide with the line of the galactic disc. The large circle indicates the location of the galactic centre.

If all GW sources were concentrated in the galactic centre, we would have found a coincidence excess twice a day. The coincidence excess occurs only once per day, just when the line of maximum sensitivity of the detectors overlaps with the galactic disc, as if GW sources were distributed in the galactic disc and not just located at its centre.

As for the energy balance, in 2001 we find in the interval 3 to 5 sidereal hours a coincidence excess of, very roughly, $n_c - \bar{n} \sim 6$ coincidences occurring in five days (two sidereal hours out of 24, in a total time period of 1490 h \sim 60 days). In terms of energy conversion into GW we have, very roughly, about one coincidence per day with a signal energy of about 100 mK. This corresponds, using the classical cross-section, to a conventional burst with amplitude $h \sim 2 \times 10^{-18}$ and to the isotropic conversion into GW energy of 0.004 solar masses, with sources located at a distance of 8 kpc. The observed rate is much larger than what the models available today predict for galactic sources. We note, however, that our rate of events is within the upper limit determined by IGEC [6] for short GW bursts and by the 40 m-LIGO prototype interferometer [16] for coalescing binary sources in the galaxy.

We think it is unlikely that the observed coincidence excess is due to noise fluctuations, but we prefer to take a conservative position and wait for a stronger confirmation of our result before reaching any definite conclusion and claiming that gravitational waves have been observed. Furthermore, although we have excluded that the events are due to cosmic ray showers (see section 6), we cannot completely rule out that they be due to some other exotic, still unknown, phenomenon. A possible way to distinguish GW from other causes is to measure other vibrational modes of the detectors and verify that, as predicted by general relativity, only quadrupole modes are excited. This requires multimode detection (with bars or spheres) and to improve the signal-to-noise ratio of the apparatus.

We expect to collect new data with EXPLORER and NAUTILUS with improved sensitivity. We plan to repeat the same analysis with these new data and also with any other new data provided by other GW groups, those which operate the resonant detectors and those which operate or are beginning to operate the interferometric detectors GEO, LIGO, TAMA and VIRGO.

Acknowledgments

We are indebted to Professor Ugo Amaldi for very useful discussions and suggestions. We thank the CERN cryogenic facility and F Campolungo, R Lenci, G Martinelli, A Martini, E Serrani, R Simonetti and F Tabacchioni for precious technical assistance.

References

- [1] Astone P *et al* 1993 *Phys. Rev. D* **47** 362
- [2] Mauceli E *et al* 1996 *Phys. Rev. D* **54** 1264
- [3] Blair D G *et al* 1995 *Phys. Rev. Lett.* **74** 1908
- [4] Astone P *et al* 1997 *Astropart. Phys.* **7** 231
- [5] Cerdonio M *et al* 1997 *Class. Quantum Grav.* **14** 1491–4
- [6] Allen Z A *et al* 2000 *Phys. Rev. Lett.* **85** 5046–50
- [7] Astone P *et al* 2001 *Class. Quantum Grav.* **18** 243–51
- [8] Astone P, Buttiglione C, Frasca S, Pallottino G V and Pizzella G 1997 *Nuovo Cimento* **20** 9
- [9] Papoulis A 1965 *Probability, Random Variables and Stochastic Processes* (New York: McGraw-Hill) p 126
- [10] Astone P *et al* 2000 *Phys. Rev. D* **62** 042001
- [11] Astone P *et al* 2002 Adaptive matched filters for the EXPLORER and NAUTILUS gravitational wave detectors (in preparation)
- [12] Weber J 1969 *Phys. Rev. Lett.* **22** 1320
- [13] Astone P, Frasca S and Pizzella G 2000 *Int. J. Mod. Phys. D* **9** 341–6
- [14] Weber J 1970 *Phys. Rev. Lett.* **25** 180
- [15] Baryshev Yu V and Paturel G 2001 *Preprint* astro-ph/0104115
- [16] Allen B *et al* 1999 *Phys. Rev. Lett.* **83** 1498